

# Congenital varicella syndrome in a mono chorionic diamniotic twin pregnancy

Vania A Villota<sup>1</sup>, Julián Delgado<sup>2</sup>, Harry Pachajoa<sup>1,3</sup>

<sup>1</sup>Centre for Research in Congenital Abnormalities and Rare Diseases, Icesi University, <sup>2</sup>Department of Gynecology and Obstetrics, Universidad del Valle, <sup>3</sup>Fundación Clínica Valle del Lili, Cali, Colombia

Congenital varicella syndrome encompasses a broad spectrum of malformations present in children of mothers who developed chickenpox during the first 20 weeks of gestation. We report a case of a mono chorionic diamniotic twin pregnancy, with maternal exposure to chickenpox during the thirteenth week of gestation, which produced one symptomatic and one healthy child.

**Key words:** Chickenpox, congenital varicella syndrome, diamniotic, herpes zoster virus, mono chorionic

**How to cite this article:** Villota VA, Delgado J, Pachajoa H. Congenital varicella syndrome in a mono chorionic diamniotic twin pregnancy. *J Res Med Sci* 2014;19:474-6.

## INTRODUCTION

Congenital varicella syndrome (CVS) is an uncommon complication of maternal varicella infection occurring in about 2% of the newborns of women infected with varicella between the seventh and the twenty-eighth weeks of pregnancy. Primary varicella infection during pregnancy can rarely result in intrauterine infection of the fetus causing congenital varicella syndrome.<sup>[1]</sup>

The typical clinical manifestations of congenital varicella syndrome include skin lesions in dermatomal distribution (73%), neurological defects (62%), ocular abnormalities (52%), limb hypoplasia (46%), and low birth weight (23%) among others minor defects.<sup>[2]</sup> We report the clinical features of the first case of congenital varicella in a mono chorionic diamniotic twin pregnancy.

Case report in the University Hospital of Valle, Cali, Colombia. 2008.

A 24-year-old woman was gravida 1, para 0, with an early uneventful twin mono chorionic diamniotic pregnancy and a tenth week ultrasound normally reported. In the thirteenth week of gestation the patient presented with fever for one week and pruritic vesicular eruptions on the skin, consistent with the classic presentation of chickenpox. The patient had no history of chickenpox in childhood and had not received immunization against it. No antiviral treatment was given. At 24 weeks of pregnancy the

patient underwent a new ultrasound in which one of the fetuses presented with decreased head circumference and facial abnormalities.

A Cesarean section was performed at 36 weeks of pregnancy. The first newborn was a 2,588 g, 45 cm long female, with Apgar scores of 2 and 0, at one and five minutes, respectively. It presented CVS stigmata, with a right hemicranial sloughing scalp, homolateral microphthalmia, microtia and hemifacial microsomia, and commissural labial deviation (Figures 1 and 2). The second newborn was a 2,750 g, 49 cm long female, with Apgar scores of 9 and 10, at one and five minutes, respectively, with no physical abnormalities.

At birth, the mother's and newborn's IgG varicella antibodies (by Enzyme-linked immunosorbent assay (ELISA)) were present at a titer of 13.3 U/mL and 9 U/mL, respectively. Both had negative specific IgM antibodies. There were no detectable antibody titers in the healthy newborn.

## DISCUSSION

There is a very high prevalence for herpes zoster virus (HZV) in the pediatric population; about 85% of people younger than 15 years of age have been exposed to the virus, and 90% of the child-bearing age women will be immune to the disease. Complications of varicella are more frequent and severe in adults than in children, and this is especially true during pregnancy, wherein, both the mother and the fetus are affected.<sup>[3]</sup> When primary



**Figure 1:** Female neonate with hemangiomas skin lesions compromising the right scalp and forehead, hemifacial microsomia, homolateral microphthalmia, microtia, and commissural labial deviation

infection with HZV occurs during gestation there is a chance of vertical transmission to the fetus at the point when the virus has colonized the mucous membranes.<sup>[4]</sup> There is a 25% risk of infection, and from these, 12% can result in CVS.<sup>[5]</sup>

The congenital varicella syndrome was first described by La Foret and Lynch, in 1947, and since then, approximately 130 cases have been reported. The estimated prevalence of CVS is one to five in 10,000 pregnancies, where primary infection in the first 20 weeks has been documented.<sup>[6]</sup> It has a 30% mortality and miscarriage rate of 3-8%.<sup>[7]</sup> The mechanism of CVS is thought to be a consequence of the reactivation of the HZV in the uterus, akin to the mechanism of HZ development, rather than being due to the primary infection.<sup>[8]</sup>

According to literature, the most common clinical features of CVS are: Skin lesions that resemble scars, burns or present with hemangiomas characteristics with a dermatomal distribution (72%); neurological defects (62%) such as cortical atrophy, microcephaly, seizures and mental retardation; ocular alterations (52%) like chorioretinitis, cataract, Horner syndrome, and nystagmus; limb hypoplasia (46%); and low birth weight (23%), among other less common features.<sup>[9]</sup>

The patient in this report presented with hemangiomas skin lesions compromising the right scalp and forehead, microphthalmia, microtia, and a right labial commissure deviation. Serological tests, clinical features, and the clear antecedent of varicella in the thirteenth week of pregnancy gave us reasons to establish a clinical and serological diagnosis. On the other hand, the lack of clinical manifestations in the other twin was remarkable, which suggested two possibilities: There was no vertical



**Figure 2:** Overview of the patient. Note that the commitment is confined exclusively to the face and skull

transmission of the infection or there was discordance in the response to exposition to the same agent. It was worth noting that the antibody titers in the fetus did not relate to the development of the syndrome, and that according to literature, there was no visible infection in 39% of the cases.<sup>[6]</sup>

In conclusion, the congenital varicella syndrome is an uncommon disease that has severe repercussions for both the mother and the fetus, which could be prevented by covering all the alternatives to prevent the infection, such as, specific and non-specific treatment and passive and active immunization, to decrease the risk of infection.<sup>[10]</sup> Thus, it must encompass a clinical assessment of the preconceptional and prenatal history in seronegative women, and exposure prevention of child-bearing age women without a history of HZV infection. In addition, further studies should be done on congenital varicella to understand the discordance among twins.

## REFERENCES

1. Seema S, Vipin S, NLS. Fetal Varicella Syndrome — A case report. *Curr Pediatr Res* 2012;16:5-8.
2. Sauerbrei A, Wutzler P. Fetales Varizellensyndrom. *Monatsschr Kinderheilkd* 2003;151:209-13.
3. Smith CK, Arvin AM. Varicella in the fetus and newborn. *Semin Fetal Neonatal Med* 2009;14:209-17.
4. Mouly F, Mirlesse V, Méritet JF, Rozenberg F, Poissonier MH, Lebon P, *et al.* Prenatal diagnosis of fetal varicella-zoster virus infection with polymerase chain reaction of amniotic fluid in 107 cases. *Am J Obstet Gynecol* 1997;177:894-8.
5. Prober CG, Gershon AA, Grose C, McCracken GH, Nelson JD. Consensus: Varicella-zoster infections in pregnancy and the perinatal period. *Pediatr Infect Dis J* 1990;9:865-9.
6. Tan MP, Koren G. Chickenpox in pregnancy: Revisited. *Reprod Toxicol* 2006;21:410-20.
7. Schulze A, Dietzsch HJ. The natural history of varicella embryopathy: A 25-year follow-up. *J Pediatr* 2000;137:871-4.

8. Lamont RF, Sobel JD, Carrington D, Mazaki-Tovi S, Kusanovic JP, Vaisbuch E, *et al.* Varicella-zoster virus (chickenpox) infection in pregnancy. *BJOG* 2011;118:1155-62.
9. Sauerbrei A, Wutzler P. The congenital varicella syndrome. *J Perinatol* 2000;20:548-54.
10. Gershon AA. Neonatal herpes simplex infection and the Three Musketeers. *N Engl J Med* 2011;365:1338-9.

**Source of support:** Nil, **Conflict of interest:** None declared.

Copyright of Journal of Research in Medical Sciences is the property of Medknow Publications & Media Pvt. Ltd. and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.